

# Three Dimensional Differential Calculus on the Quantum Group $SU_q(2)$ and Minimal Gauge Theory

D.G. Pak \*

Department of Theoretical Physics, Research Institute  
of Applied Physics, Tashkent State University,  
Vuzgorodok, 700095, Tashkent, Republic of Uzbekistan

Classification numbers: 210, 220, 240

## Abstract

Three-dimensional bicovariant differential calculus on the quantum group  $SU_q(2)$  is constructed using the approach based on global covariance under the action of the stabilizing subgroup  $U(1)$ . Explicit representations of possible  $q$ -deformed Lie algebras are obtained in terms of differential operators. The consistent gauge covariant differential calculus on  $SU_q(2)$  is uniquely defined. A non-standard Leibnitz rule is proposed for the exterior differential. Minimal gauge theory with  $SU_q(2)$  quantum group symmetry is considered.

---

\*E-mail: dmipak@apctp.kaist.ac.kr

# 1 Introduction

One of the features of non-commutative geometry in the quantum group theory [1-5] is non-uniqueness in defining a differential calculus on the quantum groups and quantum spaces. The bicovariance condition determines a unique differential calculus on the linear quantum groups  $GL_q(N)$  (up to symmetry corresponding to the exchange  $q \rightarrow \frac{1}{q}$ ) [6, 7] and provides existence of the corresponding gauge covariant differential algebra [8]. Direct reducing the  $GL_q(N)$ -bicovariant differential calculus to a case of the special linear quantum group  $SL_q(N)$  encounters difficulties connected with a loss of the centrality condition for a quantum determinant. Four-dimensional  $4D_{\pm}$  bicovariant and three-dimensional (3D) left-covariant differential calculi on the simplest special unitary quantum group  $SU_q(2)$  were considered as well using a standard Woronowicz approach [3, 6]. A full consistent construction of the 3D bicovariant differential calculus and a gauge covariant differential algebra on the  $SU_q(2)$  are unknown up to now, furthermore, there are strong limitations imposed by no-go theorems [8]. A possible way to solve this problem suggests using a non-standard Leibnitz rule as it was considered in ref. [9].

In this paper possible 3D bicovariant differential calculi and gauge covariant differential algebra on the quantum group  $SU_q(2)$  are considered in the framework of approach which respects a global  $U(1)$ -covariance. The group  $U(1)$  is a stabilizing subgroup for the quantum group  $SU_q(2)$  and the  $U(1)$ -covariant treatment allows to pass straightforward to the description of the quantum sphere  $S_q^2 \sim SU_q(2) / U(1)$ . In Section 2 we construct explicit representations of  $q$ -deformed Lie algebras of left-invariant vector fields on  $SU_q(2)$  in terms of differential operators. The  $U(1)$ -covariance constraint reduces the variety of possible covariant differential calculi on  $SU_q(2)$  and leads to a unique gauge covariant differential algebra as it is shown in Section 3. We propose non-standard Leibnitz rules for the exterior differential which are compatible with the quantum group  $SU_q(2)$  structure and gauge covariance. Section 4 is devoted to construction of the minimal quantum group gauge theory of  $SU_q(2)$ .

## 2 $q$ -deformed Lie algebras

Following the  $R$ -matrix formalism [4] the main commutation relation for the generators  $T_j^i$  ( $i, j = 1, 2$ ) of the quantum group  $SU_q(2)$  is defined by a standard  $R$ -matrix as follows

$$R_{12}T_1T_2 = T_2T_1R_{12}. \quad (2.1)$$

Let us choose a covariant parametrization for the matrix  $T_j^i$

$$T_j^i = \begin{pmatrix} y^1 & x^1 \\ y^2 & x^2 \end{pmatrix} \equiv (y^i x^i), \quad (2.2)$$

where  $x^i, y_i$  are generators (coordinates) of the function algebra on the quantum hermitean vector space  $U_q^2$  endowed with an involution  $*$  :  $x^i = y_i^*$  and  $SU_q(2)$ -comodule structure. The unimodularity condition takes a simple covariant form

$$\mathcal{D} \equiv \det_q T_j^i = x_i y^i = 1, \quad x_i = \varepsilon_{ij} x^j.$$

Hereafter the  $SU_q(2)$  indices are raised and lowered with the invariant metric  $\varepsilon_{ij}$  ( $\varepsilon_{12} = 1, \varepsilon_{21} = -\frac{1}{q}$ ).

The parametrization (2.2) was used in a harmonic formalism [10] of extended superfield supesymmetric theories. The coordinates  $(x, y)$  parametrize the quantum sphere  $S_q^2 \sim SU_q(2)/U(1)$  and are just the quantum generalizations of classical harmonic functions ( $u^\pm$ ) (so called "harmonics")

$$x^i \equiv u^{+i}, \quad y^i \equiv u^{-i}.$$

The signs  $(\pm)$  correspond to charges  $(\pm 1)$  of a stabilizing subgroup  $U(1)$  for the quantum group  $SU_q(2)$ . To simplify notations we shall not pass to the notations adopted in the harmonic formalism keeping in mind that all geometric objects (like coordinates, derivatives, differential forms etc.) have definite  $U(1)$  charges.

Consider main commutation relations between the coordinates  $(x, y)$  and derivatives  $\partial_i \equiv \frac{\partial}{\partial x^i}$ ,  $\bar{\partial}^i \equiv \frac{\partial}{\partial y_i}$  on the quantum group  $SU_q(2)$  :

$$\begin{aligned} R_{12}(\partial_T)_1(\partial_T)_2 &= (\partial_T)_2(\partial_T)_1 R_{21}, \quad (\partial_T)_j^i \equiv \begin{pmatrix} \bar{\partial}_1 & \bar{\partial}_2 \\ \partial_1 & \partial_2 \end{pmatrix}, \\ \partial_i x^k &= q^{3-2i} \delta_i^k + q Y_{mi}^{nk} x^m \partial_n, \quad \bar{\partial}^i \bar{y}_j = \delta_j^i + q y_m \bar{\partial}^n \hat{R}_{nj}^{mi}, \\ \partial_i y_j &= q(\hat{R}^{-1})_{ji}^{lk} y_k \partial_l, \quad \bar{\partial}^i x^j = \frac{1}{q} \hat{R}_{kl}^{ij} x^k \bar{\partial}^l, \end{aligned} \quad (2.3)$$

$$Y_{sj}^{ri} = (\hat{R}^{-1})_{js}^{ir} q^{2(s-i)}.$$

The commutation relations (2.3) do not differ on principle from ones given in ref. [11]. Our choice is motivated by using manifest covariant tensor notations which are convenient in constructing explicit representations for the  $q$ -Lie algebras. Thus, one implies all geometric

objects with upper (lower) indices to be transformed under the quantum group co-action  $\Delta$  like classical co-(contra-) variant tensors. For instance, a second rank tensor  $N_i^j$  will be transformed as follows

$$(N_i^j)' = (T^\dagger)_i^k T^j_l N_k^l \quad (2.4)$$

(Hereafter the signs  $\otimes$  of tensor product are omitted).

Let us define the left-invariant first-order differential operators

$$D^{++} \equiv x_i \bar{\partial}^i, \quad D^{--} \equiv -y_i \partial^i,$$

where  $(\pm\pm)$  correspond to  $U(1)$  charges  $(\pm 2)$ . The action of the operators  $D^{\pm\pm}$  on the coordinates  $(x, y)$  has a simple form

$$\begin{aligned} D^{++} x^i &= 0, & D^{--} x^i &= y^i, \\ D^{++} y_i &= x_i, & D^{--} y_i &= 0. \end{aligned}$$

The Leibnitz rule for these differential operators may be written in a convenient form if one considers their action on functions with definite  $U(1)$  charges. The functions are defined in analogy with the classical case [10] and can be decomposed in formal series

$$f^{(n \geq 0)}(x, y) = \sum_{k=1}^{\infty} C_{(i_1 i_2 \dots i_{k+n} j_1 j_2 \dots j_k)} x^{i_1} x^{i_2} \dots x^{i_{k+n}} y^{j_1} y^{j_2} \dots y^{j_k}, \quad (2.5)$$

where  $C_{(i_1 i_2 \dots i_{k+n} j_1 j_2 \dots j_k)}$  are -number coefficients symmetrized over all indices. Functions with negative charges are defined in a similar manner. After some calculations one can find the next Leibnitz rule for the operators  $D^{\pm\pm}$ :

$$D^{\pm\pm}(f^{(m)} g^{(n)}) = (D^{\pm\pm} f^{(m)}) g^{(n)} + q^{-m} f^{(m)} D^{\pm\pm} g^{(n)}. \quad (2.6)$$

This is a special feature of quantum group non-commutative geometry that the quantum analogue to classical  $U(1)$  generator can be realized as a second-order differential operator

$$D^0 \equiv -x_i \partial^i - q^2 y_i \bar{\partial}^i + (1 - q^2) x_i y_k \bar{\partial}^k \partial^i. \quad (2.7)$$

The operator  $D^0$  has eigenfunctions which are just the functions with definite  $U(1)$  charges

$$D^0 f^{(n)} = \{n\}_q f^{(n)}, \quad \{n\}_q \equiv \frac{1 - q^{-2n}}{1 - q^{-2}}, \quad (2.8)$$

where  $\{n\}_q$  is a  $q$ -number. It is not hard to check the following Leibnitz rule for the operator  $D^0$

$$D^0(f^{(m)}g^{(n)}) = (D^0f^{(m)})g^{(n)} + q^{-2m}f^{(m)}D^0g^{(n)}. \quad (2.9)$$

Reducing the space of functions on  $SU_q(2)$  to the space of functions with a definite  $U(1)$  charge one obtains the covariant description of the coset  $S_q^2 \sim SU_q(2)/U(1)$ .

By direct calculating one can verify that the operators  $D^{\pm\pm,0}$  form the  $q$ -deformed Lie algebra of  $SU_q(2)$  [12]

$$\begin{aligned} [D^0, D^{++}]_{q^{-4}} &= \{2\}_q D^{++}, & [D^0, D^{--}]_{q^4} &= \{-2\}_q D^{--}, \\ [D^{++}, D^{--}]_{q^2} &= D^0, \end{aligned} \quad (2.10)$$

here,  $[A, B]_{q^s} \equiv AB - q^s BA$ . Note, that the algebra (2.10) is valid irrespective of whether one imposes the unimodularity constraint  $\mathcal{D} = 1$ . We shall treat the algebra (2.10) as a main  $q$ -deformed Lie algebra of left-invariant vector fields on the quantum group  $SU_q(2)$ . A corresponding  $q$ -generalized Jacobi identity is available

$$\begin{aligned} [D^0, [D^{++}, D^{--}]_{q^2}] + [D^{++}, [D^{--}, D^0]_{q^{-4}}]_{q^{-2}} \\ + q^2 [D^{--}, [D^0, D^{++}]_{q^{-4}}]_{q^{-2}} = 0. \end{aligned}$$

Let us now pass to constructing other possible  $q$ -deformed Lie algebras of left-invariant vector fields on the  $SU_q(2)$ . For this purpose we consider differential operators  $\mu, \nu$  [13]

$$\mu = 1 + (q^2 - 1)y_i \bar{\partial}^i, \quad \nu = 1 + (1 - \frac{1}{q^2})x_i \partial^i. \quad (2.11)$$

One can see that the operators  $\mu, \nu$  obey the simple commutation relations

$$\begin{aligned} \mu D^{--} &= q^2 D^{--} \mu, & \mu D^{++} &= \frac{1}{q^2} D^{++} \mu, \\ \mu D^0 &= D^0 \mu, & \mu \nu &= \nu \mu. \end{aligned}$$

Similar formulae hold for the operator  $\nu$  as well. Using these relations one can find that the operators  $\mathcal{D}^{++}, \mathcal{D}^{--}, \mathcal{D}^0$  defined by the next equations

$$\begin{aligned} \mathcal{D}^{++} &= \mu^{-\frac{1}{2}} D^{++}, & \mathcal{D}^{--} &= \nu^{-\frac{1}{2}} D^{--}, \\ \mathcal{D}^0 &= \frac{1}{q} \mu \nu D^0 \equiv [\partial^0]_q \end{aligned}$$

generate just the Drinfeld-Jimbo quantum algebra

$$\begin{aligned} [\partial^0, \mathcal{D}^{++}] &= 2\mathcal{D}^{++}, & [\partial^0, \mathcal{D}^{--}] &= -2\mathcal{D}^{--}, \\ [\mathcal{D}^{++}, \mathcal{D}^{--}] &= [\partial^0]_q. \end{aligned} \quad (2.12)$$

To construct other possible  $q$ -Lie algebras one introduces another differential operators  $\Delta^{++}$ ,  $\Delta^{--}$ ,  $\Delta^0$  as follows

$$\begin{aligned}\Delta^{++} &= D^{++}, \quad \Delta^{--} = \frac{q^2 - 1}{q^{2p}} \hat{Z}^{1-p} D^{--}, \\ \Delta^0 &= \frac{1 - \hat{Z}^s}{1 - q^{2s}}, \\ \hat{Z} &\equiv (\mu\nu)^{-\frac{1}{2}}, \quad \hat{Z} f^{(n)} = q^n f^{(n)}.\end{aligned}\tag{2.13}$$

The operators  $\Delta^{\pm\pm,0}$  generate the next  $q$ -deformed Lie algebra:

$$\begin{aligned}\Delta^{++}\Delta^{--} - q^{2p}\Delta^{--}\Delta^{++} &= \frac{\hat{Z}^2 - 1}{\hat{Z}^{1+p}}, \\ \Delta^0\Delta^{++} - q^{2s}\Delta^{++}\Delta^0 &= \Delta^{++}, \\ \Delta^0\Delta^{--} - q^{-2s}\Delta^{--}\Delta^0 &= -q^{-2s}\Delta^{--},\end{aligned}$$

where  $s, p$  – arbitrary integers. The equation (2.13) allows to express the operator  $\hat{Z}$  in terms of  $\Delta^0$ , then the arbitrariness in the choice of parameters  $s, p$  can be reduced by considering only quadratic in  $\Delta^{\pm\pm,0}$   $q$ -commutators.

### 3 Gauge covariant differential algebra

In this section we give description of possible  $SU_q(2)$  bicovariant differential algebras with  $U(1)$  conserved charge. The gauge covariance condition leads to a unique differential algebra of  $SU_q(2)$ . At the same time a Leibnitz rule for the exterior differential is not fixed yet. To find the differentiation rules one needs to choose a corresponding  $q$ -Lie algebra of left-invariant vector fields.

Consider the left-invariant Cartan 1-forms  $\Omega$  on the quantum group  $SU_q(2)$

$$\Omega = dT^\dagger T \equiv \begin{pmatrix} \omega^0 & \omega^{++} \\ \omega^{--} & -q^2\omega^0 \end{pmatrix},$$

where  $\omega^0, \omega^{++}, \omega^{--}$  are the basic left-invariant differential 1-forms with corresponding  $U(1)$  charges  $(0, +2, -2)$ . One defines gauge transformations as follows

$$\begin{aligned}T^g &= \tilde{T}T, \\ \Omega^g &= \Omega - T^\dagger \tilde{\Omega} T, \quad \tilde{\Omega} \equiv d\tilde{T}^{-1} \tilde{T},\end{aligned}\tag{3.1}$$

where the matrix  $\tilde{T}$  commutes with the matrices  $T, dT$  and satisfies the same equation (2.1) as for the matrix  $T$ . In the case of gauge symmetry the matrices  $\tilde{T}, T$  depend on

the coordinates of a base space-time and the connection 1-form  $A$  has the same transformation and commutation properties as the right invariant 1-form  $dTT^\dagger$ . It turns out that the requirement of global  $U(1)$ -covariance and the consistence with the quantum group structure determine uniquely all commutation relations between the differential 1-forms  $\omega$  and the coordinates  $(x, y)$ . As a result we have

$$\begin{aligned}\omega^{++}x &= qx\omega^{++}, & \omega^{--}x &= \frac{1}{q}x\omega^{--} + \frac{1-q^4}{q}y\omega^0, \\ \omega^{++}y &= \frac{1}{q}y\omega^{++}, & \omega^{--}y &= qy\omega^{--}, \\ \omega^0x &= x\omega^0 + (1 - \frac{1}{q^2})y\omega^{++}, & \omega^0y &= y\omega^0.\end{aligned}\tag{3.2}$$

Similar consideration of commutation relations for the basic differential 1-forms  $\omega^{\pm\pm,0}$  leads to covariant algebras parametrized by a real number  $\sigma$ :

$$\omega^{++}\omega^{++} = \omega^{--}\omega^{--} = 0, \quad \omega^{\pm\pm}\omega^0 + q^{\pm 2}\omega^0\omega^{\pm\pm} = 0,\tag{3.3}$$

$$\omega^{++}\omega^{--} + q^\sigma\omega^{--}\omega^{++} + \frac{q^2(1-q^\sigma)(1+q^2)}{q^2-1}\omega^0\omega^0 = 0,\tag{3.4}$$

$$\omega^0\omega^0 = \frac{1-q^2}{q^2(1+q^2)}\omega^{++}\omega^{--}.\tag{3.5}$$

It should be noted that the algebra defined by eqs. (3.3-3.5) is bicovariant irrespective of whether one considers the last relation (3.5). Requiring the covariance under the gauge transformations and using the additional commutation constraint

$$\tilde{\Omega}\Omega = -q^2\Omega\tilde{\Omega}$$

one finds a unique gauge covariant differential algebra at  $\sigma = 4$ :

$$\begin{aligned}\omega^{++}\omega^{++} &= \omega^{--}\omega^{--} = 0, & \omega^{\pm\pm}\omega^0 + q^{\pm 2}\omega^0\omega^{\pm\pm} &= 0, \\ (1+q^2)^2\omega^0\omega^0 &= \frac{1}{q^2}\omega^{++}\omega^{--} + q^2\omega^{--}\omega^{++}.\end{aligned}\tag{3.6}$$

The equation (3.5) is not gauge covariant and should be omitted. So defined gauge covariant differential algebra differs from one considered in refs. [6,9]. Our treatment does not contain the condition of vanishing for the central element  $C_2 \equiv tr_q(\Omega^2)$ , which is not gauge covariant. Here we have used the notion of the  $q$ -deformed covariant trace [4, 7].

One can rewrite the commutation relations for the gauge covariant differential algebra in terms of the  $R$ -matrix. Direct checking leads to the next formulae

$$R_{12}dT_1T_2 = T_2dT_1R_{12},\tag{3.7}$$

$$R_{12}\Omega_2R_{12}^{-1}\Omega_1 + \frac{1}{q^2}\Omega_1R_{12}\Omega_2R_{12}^{-1} - \frac{q^2}{1+q^2+q^4}(E_{12} - (1+q^2)A_{21})tr_q\Omega^2 = 0,\tag{3.8}$$

here  $E_{kl}^{ij} = \delta_k^i \delta_l^j$  and  $A_{21}$  is the quantum antisymmetrizer [11]. Note, that the first relation in (3.7) had been obtained earlier in ref. [9].

To construct an exterior differential it is convenient to use the definition based on the dualism between the exterior algebra of differential forms and the  $q$ -Lie algebra of vector fields. In this way the Leibnitz rule is followed straightforwardly and it depends only on a special choice of the  $q$ -Lie algebra.

Let us start from a general 3D  $q$ -Lie algebra of left-invariant vector fields  $D^a = (D^{++}, D^{--}, D^0)$  on the quantum group  $SU_q(2)$  with a Lie bracket

$$[D^a, D^b]_B \equiv D^a D^b - B^{abcd} D^c D^d = C^{abc} D^c.$$

We consider the matrix  $B^{abcd}$  to be unitary, so that it generates a representation of the permutation group. Thus, one can easily define the alteration rules for the tensor algebra of vector fields. Moreover, a generalized Jacobi identity will be available as well.

The basic left-invariant differential 1-forms  $\omega^a$  are defined as dual objects by means of the scalar product  $\omega^a(D^b) = \delta^{ab}$ . The action of the exterior differential on arbitrary functions  $f$  and differential 1-forms  $u$  is defined in analogy with the classical case [14]

$$\begin{aligned} df(D^a) &= D^a f, \\ du(D^a, D^b) &= -\frac{1}{2}(D^a u(D^b) - B^{abcd} D^c u(D^d) - u([D^a, D^b]_B)), \\ du(D^a, D^b) &= -B^{abcd} du(D^c, D^d). \end{aligned} \tag{3.9}$$

Rules for the exterior differentiation of the differential ( $n > 1$ )-forms can be generalized in a similar fashion. The Cartan-Maurer equations have a standard form

$$d\omega^d(D^a, D^b) = \frac{1}{2} C^{abc} \omega^d(D^c). \tag{3.10}$$

As a concrete example we consider the  $q$ -Lie algebra (2.10) which is consistent with the gauge covariant algebra of left-invariant differential 1-forms (3.7). In that case differentiation rules (3.9) can be rewritten in a more familiar form after using the explicit tensor representation for the exterior products of 2-forms  $\omega^a \wedge \omega^b$ . After some calculations one finds

$$\begin{aligned} df &= \omega^a D^a f, \\ d(\omega^{++} f) &= d\omega^{++} f + \beta \omega^0 \omega^{++} D^0 f - d\omega^0 D^{++} f, \\ d(\omega^{--} f) &= d\omega^{--} f + \beta q^2 \omega^0 \omega^{--} D^0 f + q^2 d\omega^0 D^{--} f, \\ d(\omega^0 f) &= d\omega^0 f + \beta q^2 \omega^{++} \omega^0 D^{--} f + \beta \omega^{--} \omega^0 D^{++} f, \\ \beta &\equiv \frac{1 + q^4}{q^2(1 + q^2)}. \end{aligned} \tag{3.11}$$



It should be noted that the formulae (3.11) involve just three independent basis differential 2-forms  $\omega^0\omega^{++}, \omega^0\omega^{--}, d\omega^0$  in the space of exterior 2-forms in correspondence with the classical case. The fourth linearly independent basis 2-form  $\sigma^0$  can be defined as follows

$$\sigma^0 = \frac{1}{1+q^2}(\omega^{++}\omega^{--} + q^2\omega^{--}\omega^{++}),$$

The form  $\sigma^0$  takes a non-zero value only for the symmetrical tensor product  $D^0 \otimes D^0$ :

$$\sigma^0(D^0, D^0) = \rho,$$

where the number  $\rho$  vanishes in the classical limit  $q \rightarrow 1$ . Due to this property the form  $\sigma^0$  does not appear in eqs. (3.11).

To construct the differentiation rules for the  $(n > 1)$ -forms it is convenient to use the specific structure of the exterior algebra of  $SU_q(2)$ . It is easy to check that the invariant 2-form

$$C \equiv \frac{\beta q^4}{1+q^2+q^4} C_2 = \frac{\beta q^2}{1+q^2}(\omega^{++}\omega^{--} + \omega^{--}\omega^{++}) \quad (3.12)$$

is a central element. We put the natural constraint

$$d(C\omega^{(n)}f) = Cd(\omega^{(n)}f). \quad (3.13)$$

Using this constraint and starting from the most general form for differentiation rules one finds

$$\begin{aligned} d(\omega^0\omega^{\pm\pm}f) &= \pm q^{\mp 4}vD^{\pm\pm}f, \\ d(\omega^{++}\omega^{--}f) &= \frac{1}{\beta}Cdf + vD^0f, \\ d(\omega^{--}\omega^{++}f) &= \frac{1}{\beta q^2}Cdf - vD^0f, \\ d(\omega^0\omega^{++}\omega^{--}f) &= \frac{1}{\beta}Cd\omega^0f - q^4C\omega^0\omega^{++}D^{--}f - \frac{1}{q^2}C\omega^0\omega^{--}D^{++}f, \\ d(vf) &= 0, \\ v &\equiv \frac{1}{2}(\omega^0\omega^{++}\omega^{--} - q^2\omega^0\omega^{--}\omega^{++}), \end{aligned} \quad (3.14)$$

where  $v$  is a volume 3-form on the  $SU_q(2)$ .

All basis differential forms of order  $n > 3$  can be obtained from lower order forms multiplied by the invariant  $C$  in an appropriate degree. So that the relations (3.11, 3.14) complete the differential rules for the differential algebra of  $SU_q(2)$ .

Having carried out some calculations one can also find the explicit expressions for the Cartan-Maurer equations (3.10)

$$d\Omega = \Omega^2 - \frac{q^2}{1+q^2}I \operatorname{tr}_q \Omega^2.$$

The right hand side of the equation contains only the traceless part of  $\Omega^2$ .

## 4 Construction of the minimal $SU_q(2)$ gauge theory

A minimal gauge theory corresponding to the quantum algebra  $SU_q(2)$  was proposed in ref. [15], where the initial gauge transformations contain by definition the antipodal map. One can try to formulate a gauge Yang-Mills theory for the quantum group  $SU_q(2)$  in analogy with a covariant  $GL_q(N)$  version proposed in ref. [7]. For this purpose one should define the algebra of main operators (matter fields and gauge potential) and a corresponding comodule structure.

In the case of the quantum group  $SU_q(2)$  it is convenient to put the matter field  $\phi^i(\bar{\phi}^i)$  into one matrix  $\Phi = (\bar{\phi}^i \ \phi^i)$ . The operators  $(\Phi, d\Phi)$  generate the  $Z_2$ -graded algebra  $Z$  with the same commutation relations as for the differential algebra of  $SU_q(2)$ . The algebra  $Z$  is a left  $SU_q(2)$ -comodule with the following co-action:

$$\Phi \rightarrow \Phi^g = T\Phi, \quad (4.1)$$

$$d\Phi \rightarrow (d\Phi)^g = (dT)\Phi + T(d\Phi). \quad (4.2)$$

All axioms for the comodule are fulfilled.

We introduce also an operator  $A$  (gauge potential 1-form) satisfying the same commutation relations as for the right-invariant Cartan 1-forms on the  $SU_q(2)$ . One can consider the quantum analogue to the gauge transformation for the  $A$ :

$$A \rightarrow A^g = TAT^\dagger + (dT)T^\dagger. \quad (4.3)$$

A covariant differential  $\Delta$  acting on the matter field is defined as

$$\Delta\Phi = (d - A)\Phi. \quad (4.4)$$

The curvature 2-form  $F$  is introduced as follows

$$F = dA - A^2 + \frac{1}{1+q^2} tr_q A^2 \quad (4.5)$$

and it contains only traceless part.

To define the commutation relations in the algebra  $G$  generated by the operators  $(\Phi, d\Phi, A, dA)$  we will take into account the compatibility conditions with the gauge transformations and the centrality property for the quantum determinant  $\det_q \Phi = 1$ . The commutation relations for the operators  $\Phi, A$  are uniquely defined from the formula (3.7), since the gauge potential  $A$  in a pure gauge limit is just the right-invariant Cartan form on the quantum group  $SU_q(2)$ , so we have

$$\Phi_1 A_2 = R_{21} A_2 R_{21}^{-1} \Phi_1. \quad (4.6)$$

We demand the covariant combination  $(F\Phi)$  to have the same commutation relations with itself and  $A$  as for the operator  $\Phi$

$$(F\Phi)_1 A_2 = R_{21} A_2 R_{21}^{-1} (F\Phi)_1, \quad (4.7)$$

$$R_{12} (F\Phi)_1 (F\Phi)_2 = (F\Phi)_2 (F\Phi)_1 R_{12}. \quad (4.8)$$

Using these formulae one can easily derive the next commutation relations:

$$F_1 R_{21} A_2 R_{21}^{-1} = R_{21} A_2 R_{21}^{-1} F_1, F_1 R_{21} \tilde{r}_2 R_{21}^{-1} = R_{21} \tilde{r}_2 R_{21}^{-1} F_1, \quad (4.9)$$

where  $\tilde{r} = (d\Phi)\Phi^\dagger$ .

The bicovariance condition, the traceless property for the  $F$  and the centrality of the quantum determinant  $\det_q \Phi$  lead to the following commutation relation for the operators  $(\Phi, F)$ :

$$\Phi_1 F_2 = R_{21} F_2 R_{21}^{-1} \Phi_1. \quad (4.10)$$

From this equation taking into account the eqn. (34) one finds

$$R_{12} F_1 R_{21} F_2 = F_2 R_{12} F_1 R_{21}. \quad (4.11)$$

Note, that the last equation coincides with the corresponding relation in  $GL_q(N)$  gauge theory [7]. It is not difficult to check the next simple relations as well:

$$R_{12} \Delta \Phi_1 \Phi_2 = \Phi_2 \Delta \Phi_1 R_{12}, \quad (4.12)$$

$$\Phi_1 dA_2 = R_{21} dA_2 R_{21}^{-1} \Phi_1. \quad (4.13)$$

The commutation relations obtained above are the main ones which imply all other commutations in the algebra  $G$ . To construct a formal expression for the Lagrangian of the gauge theory one needs to specify the underlying space-time and define the dual  $*$ -operation. The simplest variant correspond to the choice of the space-time isomorphic to the quantum space of  $SU_q(2)$ . Another problem is the construction of Leibnitz rules for the exterior differential which have essentially non-standard form. Differentiation rules for the algebra of matter fields  $(\Phi^i, d\Phi)$  are defined by similar formulae (23, 26) in full analogy with the case of the differential algebra of  $SU_q(2)$ . For instance, one has the following equations for the quadratic combinations of matter fields

$$\begin{aligned} d(\phi^i \phi^j) &= d\phi^i \phi^j + \frac{1}{q^2} \phi^i d\phi^j + \frac{q^2 - 1}{q^3} \varepsilon^{ij} d\phi_k \phi^k, \\ d(\bar{\phi}^i \bar{\phi}^j) &= d\bar{\phi}^i \bar{\phi}^j + q^2 \bar{\phi}^i d\bar{\phi}^j, \\ d(\bar{\phi}^i \phi^j) &= d\bar{\phi}^i \phi^j + q^2 \bar{\phi}^i d\phi^j + (q^2 - 1) \bar{\phi}^i \bar{\phi}^j d\phi_k \phi^k. \end{aligned}$$

Another possible way towards a consistent minimal quantum group gauge Yang-Mills theory corresponds to the differential calculus with a  $q$ -Lie algebra differed from one defined by eqs. (2.10).

## Acknowledgments

The author would like to acknowledge Professor M. Arik and the members of the Organizing Committee for kind hospitality and financial support. Author thanks F. Mueller-Hoissen, V.D. Gershun and M. Nomura for useful discussions.

## References

- [1] V.G. Drinfeld *Quantum Groups*, in: Proc. Int. Cong. Math. **1** (Berkeley, CA, USA, 1986), 793.
- [2] M. Jimbo, Int. J. Mod. Phys. **A4** (1989) 3759.
- [3] S.L. Woronowicz, Publ. Res. Inst. Math. Sci., Kyoto University **23** (1987) 117.
- [4] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Algebra i Analiz. **1** (1989) 178.
- [5] S.L. Woronowicz, Comm. Math. Phys. **122** (1989) 125.
- [6] A.P. Isaev and P.N. Pyatov, Phys. Lett. **A179** (1993) 81.
- [7] A.P. Isaev and Z. Popowicz, Phys.Lett. **B307** (1993) 353.
- [8] I.Ya.Aref'eva and G.E.Arutyunov, "On \*-representations of the  $Z_2$ -graded extension of the quantum group  $U_q(2)$ " Trudi MIAN, v.203 (1994)
- [9] L.D. Faddeev and P.N. Pyatov, *The Differential Calculus on Quantum Linear groups*. Preprint hep-th/9402070 (1994).
- [10] A. Galperin, E. Ivanov, S. Kalitzin, V. Ogievetsky and E. Sokatchev, Class. Quant. Grav. **1** (1984) 469.
- [11] J. Wess and B. Zumino, Nucl. Phys. (Proc. Suppl.) **B18** (1990) 302.
- [12] A. Schirrmacher, J. Wess and B. Zumino, Zeit. Ph. **C49** (1991) 317.
- [13] O. Ogievetsky, Lett. Math. Phys. **24** (1992) 245.

- [14] S. Kobayashi and K. Nomizu, Vol.1. *Foundations of Differential Geometry* (Interscience Publishers, N.Y., London, 1963).
- [15] T. Sudbery Phys.Lett. B375 (1996) 75.